




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Subclinical myocardial injury, coagulopathy, and inflammation in COVID-19:
A meta-analysis of 41,013 hospitalized patients

Oluwabunmi Ogungbe, Baridosia Kumbe, Oluwadamilola Fadodun, T. Latha,
Diane Meyer, Adetoun Asala, Patricia M. Davidson, Cheryl R. Dennison
Himmelfarb, Wendy S. Post, Yvonne Commodore-Mensah

PII: S2352-9067(21)00238-4
DOI: <https://doi.org/10.1016/j.ijcha.2021.100950>
Reference: IJCHA 100950



To appear in: *IJC Heart & Vasculature*

Received Date: 23 October 2021
Revised Date: 8 December 2021
Accepted Date: 29 December 2021

Please cite this article as: O. Ogungbe, B. Kumbe, O. Fadodun, T. Latha, D. Meyer, A. Asala, P.M. Davidson, C.R. Dennison Himmelfarb, W.S. Post, Y. Commodore-Mensah, Subclinical myocardial injury, coagulopathy, and inflammation in COVID-19: A meta-analysis of 41,013 hospitalized patients, *IJC Heart & Vasculature* (2021), doi: <https://doi.org/10.1016/j.ijcha.2021.100950>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Subclinical myocardial injury, coagulopathy, and inflammation in COVID-19: A meta-analysis of 41,013 hospitalized patients

Authors:

Oluwabunmi Ogungbe, MPH, RN^{1*}
Baridosia Kumbe, BSc¹
Oluwadamilola Fadodun, BNSc²
Latha T., PhD, MSc³
Diane Meyer, MPH, RN¹
Adetoun Asala, MPH⁴
Patricia M. Davidson, PhD, MED, RN^{1,5}
Cheryl R. Dennison Himmelfarb PhD, RN^{1,6,7}
Wendy S. Post, MD, MS^{6,7}
Yvonne Commodore-Mensah, PHD, MHS, RN^{1,6,7}

Affiliations

¹Johns Hopkins University School of Nursing, Baltimore, MD, USA.

²The University of Lethbridge, Lethbridge, Alberta, Canada.

³Manipal Academy of Higher Education, Manipal, India.

⁴Jackson State University, Jackson, MS, USA.

⁵University of Wollongong, Wollongong, Australia.

⁶Johns Hopkins University School of Medicine, Baltimore, MD, USA.

⁷Johns Hopkins University Bloomberg School of Public Health, Baltimore, MD, USA.

*Corresponding Author

Oluwabunmi Ogungbe, MPH, RN¹
Johns Hopkins School of Nursing
525 N. Wolfe St.
Baltimore, MD. 21205
Email: oogungb3@jh.edu

PROSPERO database registration number: CRD42021229341

Word count: 2,995 (excluding abstract, references, tables and figure legends)

Tables: 1

Figures: 3

Abstract

Background: Infection with the SARS-CoV-2 virus can lead to myocardial injury, evidenced by increases in specific biomarkers and imaging.

Objective: To quantify the association between biomarkers of myocardial injury, coagulation, and severe COVID-19 and death in hospitalized patients.

Methods: Studies were identified through a systematic search of indexed articles in PubMed, Embase, CINAHL, Cochrane, Web of Science, and Scopus, published between December 2019 to August 2021. Effect estimates from individual studies for association between markers of myocardial injury (Troponin), myocardial stretch (N-terminal-pro hormone BNP, NT-proBNP), and coagulopathy (D-Dimer) and death or severe/critical COVID-19 were pooled using inverse variance weighted random-effects model. Odds Ratios (OR), Hazard Ratios (HR), and 95% Confidence Intervals (CI) were pooled separately and reported by outcomes of critical/severe COVID-19 and death. A meta-analysis of proportions was also performed to summarize the pooled prevalence of co-morbidities in patients hospitalized with COVID-19.

Results: We included 62 articles, with a total of 41,013 patients. The pooled proportion of patients with history of hypertension was 39% (95% CI: 34-44%); diabetes, 21% (95% CI: 18%-24%); coronary artery disease, 13% (95% CI: 10-16%); chronic obstructive pulmonary disease, 7% (95% CI: 5-8%), and history of cancer, 5% (95% CI: 4-7%). Elevated troponin was associated with higher pooled odds of critical/severe COVID-19 and death [Odds Ratio (OR: 1.76, 95% CI: 1.42-2.16)]; and also separately for death (OR: 1.72, 95% CI: 1.32-2.25), and critical/severe COVID-19 (OR: 1.93, 95% CI: 1.45-2.40). Elevations in NT-proBNP were also associated with higher severe COVID-19 and death (OR: 3.00, 95% CI: 1.58-5.70). Increases in D-dimer levels was also significantly associated with critical/severe COVID-19 and death (pooled OR: 1.38, 95% CI: 1.07-1.79).

Conclusions: This meta-analysis synthesizes existing evidence showing that myocardial injury, and coagulopathy are complications of COVID-19. The durability of these complications and their contributions to long-term cardiac implications of the disease is still being investigated. Patients who have recovered from COVID-19 may benefit from minimally invasive assessment for markers of myocardial injury, stretch and coagulopathy for early risk stratification purposes.

Keywords: COVID-19, SARS-CoV-2, Myocardial Injury, Inflammation, Troponin, Meta-analysis

Background

While COVID-19 may begin as a respiratory disease, sometimes causing pneumonitis and severe acute respiratory distress syndrome (ARDS), other major organs may also be affected, particularly the cardiovascular system, kidneys, and brain.¹ COVID-19 is associated with a wide range of symptoms, and a spectrum of clinical illnesses characteristic of multisystem disease with cardiovascular consequences.^{2,3} The characteristic of immune system activation may result in cytokine storm with the release of massive amounts of cytokines, causing both local and systemic inflammatory response, which contributes to disarray in immune, cardiac and inflammatory biomarkers.^{4,5}

Infection with SARS-CoV-2 virus can lead to elevations in cardiac biomarkers and electrocardiographic changes that are associated with poorer clinical outcomes, including ICU admission, mechanical ventilation and death.⁶⁻⁸ These cardiac biomarkers include troponin (cTnI), n-terminal pro-b-type natriuretic peptide (NT-proBNP), galectin-3, d-dimer, interleukin-6 (IL-6), and c-reactive protein (CRP).⁹⁻¹¹ In one study of 2,736 patients hospitalized for COVID-19 in New York City, small elevations in cTnI laboratory values were associated with increased mortality. NT-proBNP, a hormone that is predominantly produced by ventricular myocytes in response to increased ventricular wall stress, is an established biomarker for diagnosis and prognosis of heart failure.¹² Elevated NT-proBNP is associated with increased risk of cardiogenic shock,¹³ acute heart failure,^{14,15} myocardial infarction,¹⁴ right ventricular dysfunctions, left ventricular dysfunction,^{16,17} stress cardiomyopathy,^{16,17} arrhythmias,¹³⁻¹⁵ and venous and arterial thrombosis^{18,19} among hospitalized COVID-19 patients²⁰. According to the American College of Cardiology (ACC), based on case reports from China, US, and Europe, about 40% of hospitalized COVID-19 patients have cardiovascular or cerebrovascular disease, 16.7% of them developed arrhythmias, and 7.2% developed acute cardiac injury.²¹⁻²³

Understanding accumulating evidence on the cardiac consequences of infection with SARS-CoV-2 virus is critical to better understanding biomarkers that may indicate poorer clinical outcomes. Thus, we conducted a systematic review and meta-analysis to quantify the association between biomarkers of myocardial injury (troponin), myocardial stretch (NT-proBNP, Gal-3), coagulation (D-Dimer), inflammation (CRP, IL-6) and severe COVID-19 or death in hospitalized patients. We also calculated pooled proportions of underlying co-morbidities (hypertension, CAD, diabetes, COPD, and cancer) from the articles reviewed.

Methods

Search Strategy

This systematic review and meta-analysis followed recommendations from Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (**eTable 1 in the Supplement**). A detailed literature search was conducted in relevant scientific databases, including MEDLINE (PubMed), Embase, CINAHL, Cochrane, Web of Science, and Scopus. Search terms and strategy were developed with the help of an informationist to search for studies conducted between December 2019 and January 2021. An updated search was conducted for articles between January and August 2021. The complete search strategy for each database is available in **eTable 2 in the Supplement**. This review was registered in the PROSPERO database with registration number CRD42021229341.

Study selection and eligibility criteria

All studies identified via databases searches were imported into Covidence.²⁴ Three reviewers (O.O., B.K., O.A.) independently screened identified studies for title and abstract eligibility. Discrepancies and divergencies were discussed and adjudicated with the help of two independent reviewers (L.T. and A.A.). The following criteria full-text review inclusion were: 1) study type: retrospective, prospective, observational, or case-control studies, reporting on cardiac biomarkers (specifically Troponin I or T, NT-proBNP, CRP, IL-6, and Gal-3); 2) population: adult patients diagnosed with COVID-19; 3) exposure/intervention: confirmed COVID 19 infection; 4) outcome

indicators: myocardial injury, severe or critical clinical status, and death. Studies excluded include case reports, editorials, letters to the editor and correspondence, and conference abstracts that did not contain primary data. Preprints and non-peer reviewed articles, other review articles, and articles not published in English were also excluded.

Data extraction

Following title/abstract screening, three reviewers (O.O., B.K., O.A.) worked in pairs to independently assess the studies for full-text eligibility. A third and fourth reviewer performed conflict resolution, one adjudicator per article, to reach a consensus (L.T., and A.A.). After full-text eligibility screening, data were extracted independently by a total of five reviewers (O.O., B.K., O.A., L.T., and D.M); for each article, there were two primary independent reviewers and one adjudicator. Discrepancies for extracted data were resolved by consensus, results adjudicated where applicable and confirmation checks were done. A predesigned and piloted data template in Covidence was used for data extraction. Data extracted included author, publication year, study design, sample size, participant characteristics, proportion of comorbidities in the sampled population, median or mean levels of cardiac biomarkers and their cut-off points, outcomes of interest (COVID-19 severity, death, ICU admission, etc.), and reported measures of association.

Definitions

For this review, data on severe COVID-19 was abstracted based on the definition and categories as reported in the reviewed articles. Studies conducted in China defined severity of COVID-19 according to the Chinese guideline for COVID-19 management;²⁵ severe cases were defined as the presence of at least one of the following: (i) respiratory rate >30 breaths per minute; (ii) oxygen saturation (SpO_2) $\leq 93\%$; and (iii) PaO_2/FiO_2 ratio ≤ 300 mmHg. Critical cases were defined as those including at least one of the following: (i) respiratory failure requiring mechanical ventilation; (ii) shock; (iii) presence of other organ damage apart from respiratory; and (iv) admission to intensive care unit. Other studies used the World Health Organization definitions of severe pneumonia (Adolescent or adult: fever or suspected respiratory infection, plus one of respiratory rate >30 breaths/min, severe respiratory distress, or $SpO_2 < 90\%$ on room air) or ARDS (Onset: new or worsening respiratory symptoms within one week of known clinical insult; Chest imaging (radiograph, CT scan, or lung ultrasound): bilateral opacities, not fully explained by effusions, lobar or lung collapse, or nodules; Origin of edema: respiratory failure not fully explained by cardiac failure or fluid overload; o exclude hydrostatic cause of edema if no risk factor present; Oxygenation (adults): $PaO_2/FiO_2 \leq 100$ mmHg with PEEP ≥ 5 cmH₂O, or non-ventilated).²⁶ Majority of the articles defined myocardial injury as hs-Troponin I (hs-TnI) ≥ 99 th percentile of the upper reference limit, regardless of new abnormalities in electrocardiography and echocardiography.²⁷

Quality assessment

Quality Assessment was conducted using the Newcastle Ottawa Quality Assessment Scale (NOS)²⁸ which assesses design quality of non-randomized, cohort and case-control studies. A maximum score of 9 was aggregated across three domains of the NOS scale: selection of study groups (4 possible points), comparability of groups (2 possible points), and ascertainment of outcomes (3 possible points). The overall risk of bias was classified as either “high”, “some concern”, or “low” based on the cumulative NOS scores. The risk-of-bias assessment summary table and plots were created using the *robvis* application.²⁹

Data synthesis and statistical analysis

Analysis was conducted using the *meta* commands in Stata/IC 16.1.³⁰ All reported medians or mean values for biomarkers: Troponin I or T, NT-proBNP, Gal-3, IL-6, D-Dimer, and CRP were summarized in the original unit measured in the studies. When studies did not report mean and standard deviations, reported sample sizes, median and interquartile ranges were abstracted. Measures of associations reported in relevant studies (odds or hazard ratios) and the primary outcomes were summarized. For the meta-analysis, unadjusted hazard ratios (HR) and odds ratios (OR) and their 95% confidence intervals

(CI), reported from each study were natural log-transformed.³¹ OR, HRs and 95% CIs were pooled separately and reported by outcomes of death or critical/severe COVID-19. Random effect models with the inverse variance method were used to derive pooled measures of association, and the 95% CI and p-values to quantify the associations between elevated biomarkers and cardiovascular complications, and COVID-19 severity and death.

The pooled effect sizes were represented as forest plots. Heterogeneity of study estimates was assessed by Hedge's I^2 statistic, which was interpreted as $I^2 < 25\%$, 25-50%, 50-75%, and $> 75\%$, which indicates no, low, moderate, and high between-study heterogeneity.³² The risk of publication bias was assessed using funnel plots. Egger's meta-regression was performed to assess for small-study effects. We visually inspected the distribution of relevant studies for asymmetry or by p-value < 0.1 , indicating possible publication bias.³³

To calculate pooled proportions from the reported underlying co-morbidities (hypertension, CAD, diabetes, COPD, and cancer) from each article, we performed a meta-analysis of proportions of this binomial data, using the *metaprop* package in Stata.³⁴ Using a random-effects model, we calculated 95% exact confidence intervals for each study and the overall pooled estimate for each comorbidity. We reported the measure of heterogeneity (I^2) and displayed the outputs using forest plots.

Results

Figure 1 shows the PRISMA flow chart for study selection. A total of 3,106 records were identified through the databases on initial search, and 2,032 records from the updated search; 2,454 duplicates were removed, and 2,684 titles and abstracts were screened for inclusion. Following title and abstract screening, 2,274 articles were excluded leaving 410 articles for full-text retrieval, 58 of which full-text versions were unretrievable or conference abstracts. Upon screening, a total of 290 articles were further excluded, leaving 62 articles that met the inclusion criteria and were included in the review and meta-analysis for this study. The PRISMA flowchart was completed according to the updated 2020 guideline,³⁵ and generated using an available online Shiny App.³⁶

Study Characteristics

The characteristics of the 62 studies included are presented in **Table 1**. Most studies were conducted in China (50%), Italy (14%), the United States (12%), Turkey (3%), Spain (3%), the United Kingdom (2%), France (2%), Iran (2%), Korea (2%), Mexico (2%), Morocco (2%), Brazil (2%), Pakistan (2%), and the Netherlands (2%). Total sample size from the studies was 41,013, ranging from 38 to 11,159 patients, and median age of participants ranged from 44.6 to 75 years. The pooled proportion of participants who had severe COVID-19 or died during hospitalization was 16.8% (1,978/11,764).

Study quality

For the papers included in the meta-analysis, 24 were rated to have a low risk of bias, while 16 were rated to have a high risk of bias. Four were rated as unclear (**eFigure 1 in the Supplement**). Those studies assessed to have a high risk of bias were primarily due to incomplete outcome reporting.

Underlying cardiovascular diseases and comorbidities

A majority of the studies reported history of comorbidities among patients included in the study. The pooled proportion of patients with hypertension was 39% (95% CI: 35-43%), ranging from 15-73% (**Figure 2**). For history of diabetes, the pooled proportion was 21% (18%-23%), this ranged from 8%-44%. The pooled proportion for history of CAD was 13% (95% CI: 11-16%), and this ranged from 1-49%. In the total sample, the pooled proportion of patients who had COPD was 7% (95% CI: 6-9%), which ranged from 2-23%, for history of cancer, the pooled proportion was 6% (95% CI: 5-7%) which also ranged from 2-16% (**eFigures 4-7 in the Supplement**).

Biomarkers of myocardial injury, stretch, and coagulation

For studies that reported myocardial injury, the proportion of patients who developed myocardial injury was 18.5% (3,943/ 21,367) (**eTable 3 in the Supplement**). The overall pooled odds ratio for the association between elevated troponin levels and death and critical/severe disease was 1.76 (95% CI: 1.42, 2.16); for the outcome of death, the pooled odds was OR: 1.72 (95% CI: 1.32, 2.25); and for the outcome of critical/severe COVID-19, the pooled odds ratio was 1.93 (95% CI: 1.45, 2.40) (**Figure 3**). Similarly, the overall hazards ratio for elevated troponin and death and critical/severe disease was 1.55 (95% CI: 1.36, 1.77). The hazards for elevated troponin and death was 1.51 (95% CI 1.31, 1.75), and for critical or severe COVID-19 was 1.75 (95%CI 1.48, 2.10).

Elevations in NT-proBNP were also associated with severe COVID-19 and death: OR: 3.00 (95% CI: 1.58-5.70). The hazards ratio associated with elevated NT-proBNP and death, or critical/severe COVID-19 was 1.65 (pooled HR: 1.65, 95%CI: 0.88-3.10) (**Figure 3**). Increased D-dimer levels were also significantly associated with death and severe disease, pooled OR: 1.38 (95%CI: 1.07-1.79). While the pooled hazards ratio associated with either death, critical/severe disease, and elevated D-dimer levels was HR: 1.16 (95% CI: 0.9, 1.48) (**Figure 3**).

Abnormal increases in CRP were also associated with higher severe COVID-19 and death: OR: 1.60 (95% CI: 1.13-2.25). The hazards ratio associated with increased CRP and death or critical/severe COVID-19 was 1.15 (pooled HR: 1.15, 95% CI: 0.94-1.42) (**eFigure 2 in Supplement**). Elevated IL-6 levels were also significantly associated with death and severe disease, pooled OR: 1.55 (95% CI: 1.14-2.14) (**eFigure 3 in the Supplement**).

Discussion

In this meta-analysis, we aimed to assess the association between biomarkers of myocardial injury and stretch, coagulation, and severe COVID-19 and death. From 62 articles reviewed, hospitalized COVID-19 patients had a high prevalence of underlying comorbidities such as hypertension, diabetes, COPD, CAD, and malignancy. Furthermore, the pooled data demonstrated that elevated biomarkers of inflammation, subclinical myocardial injury, coagulation, were significantly associated with severe COVID-19 and death.

Our findings are consistent with previous findings, which have reported early in the COVID-19 pandemic that individuals who have an underlying or preexisting cardiovascular disease such as heart failure, coronary heart disease and risk factors such as being of older age, hypertension, diabetes are over-represented in COVID-19 hospitalization.³⁷ In similar meta-analyses examining the association between COVID-19 and poor COVID-19 outcomes, there is a strong association between biomarker evidence of cardiac injury and worse COVID-19 outcomes; they include elevation of cTn, NT-proBNP, d-dimer, which predict poor clinical outcomes.^{38, 39}

Cardiac involvement in COVID-19 is determined by the extent of the viral inoculum, and magnitude of host immune response, and the presence of underlying comorbidities.¹⁰ Some of the mechanisms through which direct and indirect cardiac injury may occur in the context of COVID-19 are through inflammation, endothelial activation, and microvascular thrombosis (**eFigure 8 in the Supplement**).^{10, 40} In direct viral myocardial invasion, the outer membrane spike of the SARS-CoV-2 virus has a high affinity for the ACE2 receptors and the protease transmembrane protease serine 2 (TMPRSS2), which are highly expressed in cardiac tissues.⁴¹ Hence, direct viral myocardial invasion is highly plausible, and evidence of SARS-CoV-2 positivity in cardiac tissues has been documented in autopsy reports.⁴² The activation of macrophages (major sources of cytokines and the inflammatory cytokine tumor necrosis factor (TNF)- α) is promulgated by metalloproteinase domain 17 (ADAM-17), which is also responsible for shedding of ACE2. Loss of ACE2 receptor density due to binding from the SARS-CoV-2 spike protein leads to accumulation of Angiotensin II (Ang II), and continued triggering of ADAM-17.⁴³ This creates vicious positive feedback of activated ADA-19, more ACE2 shedding, and increased Ang II-mediated inflammatory responses, which is partially responsible for the cytokine storm characteristic of the SARS-CoV-2 immune response.^{10, 44}

In the context of COVID-19, plaque destabilization and eruption can be facilitated by viral products from the systemic circulation, which could activate immune receptors on cells in already existing plaques in coronary vasculature.⁴⁵ The ongoing infection and inflammation could also lead to dysregulation of coronary vascular endothelial function leading to vasoconstriction and thrombosis.⁴⁶ Endothelial dysfunction is expressed through alteration in the vessel barrier, promotion of a coagulative state, induction of endothelial inflammation, and mediation of leukocyte infiltration.⁴⁷ Importantly, myocardial oxygen supply and demand may result from the following: endothelial dysfunction in coronary microcirculation; fixed coronary atherosclerosis limiting myocardial perfusion, high levels of circulating Ang II and arteriolar vasoconstriction resulting in systemic hypertension, and hypoxemia from ARDS or pulmonary thrombosis.¹⁰ The immense physiologic demands that result from the SARS-CoV-2 infection response and the systemic inflammatory cascade, may be sufficient to trigger this supply-demand mismatch, even in the absence of an atherothrombotic plaque.¹⁰ Another possible contributory mechanism associated with high fatality in COVID-19 and complications is disseminated intravascular coagulation (DIC), with the cross-talk between inflammation and coagulation mediated through protease-activated receptors (PARS); there have been recommendations for clinical use of direct oral anticoagulants to inhibit PARS in acute care of COVID-19 and in patients experiencing persisting symptoms of COVID-19.⁴⁸

Evidence is emerging on the long-term cardiac sequela of COVID-19-related myocardial injury, with evidence of myocardial fibrosis or myocarditis in 9-78% of patients who have had COVID-19.³⁷ Ongoing inflammation has been reported in 60%, and cardiac involvement in 78% of patients recovered from acute COVID-19.⁴⁹ Based on the results of our meta-analysis and other similar studies, and in light of post-acute sequelae of COVID-19, it is important to consider measurements of sustained expression of biomarkers signaling inflammation and myocardial damage in persons who have previously tested positive for COVID-19. Such sustained expression of certain inflammatory biomarkers are yet to be characterized in COVID-19; however, these may indicate long-term hyperinflammatory state with cardiac involvement from damage mediated by the virus during the acute phase, or even possibilities of host viral reservoir, which is not uncommon in other viruses, but yet to be established in COVID-19.⁵⁰ Considerations of the prognostic value of these biomarkers in persons previously hospitalized for COVID-19, and identification of subclinical myocardial injury can help with risk stratification, and downstream decisions about care.

Strengths and Limitations

This study has some limitations. First, we could not find a sound and acceptable method for pooling reported medians and interquartile ranges in the literature. This was challenging considering that biomarker values are not typically normally distributed, hence cannot be confidently pooled without introducing bias, particularly when using meta-analysis methods that are reliant on means and standard deviation as measures of central tendency and dispersion. Second, several of the biomarkers were reported in different units, some of which could not be confidently converted to a single unit for use in statistical analyses. This severely limited our inclusion of the biomarker values reported in the articles in the meta-analyses. Third, this review was limited to articles published in English; excluding non-English articles during screening may have introduced some bias.

Nevertheless, this study has major strengths. The meta-analytic approach combined both observational and comparative methods. This included a meta-analysis of proportions for estimating the pooled prevalence of underlying comorbidities, and a comparative meta-analysis of odds and hazards of the associations between elevated biomarker levels and severe COVID-19 and/or death. In addition, our study comprehensively examined biomarkers that are representative of the different pathways of immune response and cardiac injury associated with the COVID-19. This study also offers an updated summary of existing evidence on the relationship between biomarkers of inflammations, myocardial injury in COVID-19, and their value in prognosis. Our meta-analysis included 41,013 patients from diverse global regions and contexts. Findings from our study also corroborate with findings from other studies showing an over-representation of persons with pre-existing comorbidities in persons with poor COVID-19 outcomes.

Conclusions

Our systematic review and meta-analysis showed significant associations between markers of myocardial injury and stretch, and coagulopathy with poor COVID-19 outcomes. There is also evidence in the literature of persisting symptoms suggestive of complications in patients recovered from COVID-19. The durability of these complications and their contributions to long-term cardiac implications of the disease is still being investigated. Recovered patients, may benefit from minimally invasive assessment for markers of myocardial injury and stretch and coagulopathy for early risk stratification purposes.

Acknowledgments

We would like to acknowledge the information specialist, Stella Seal; Williams H. Welch Medical Library, Johns Hopkins Medicine, who curated the search strategy and performed the initial literature search. We would also like to acknowledge Dr. Chakra Budthrodaki, Johns Hopkins University, who reviewed the biostatistics and analytical methods used in this meta-analysis.

Disclosures

The authors report no relationships that could be construed as a conflict of interest.

Funding Source

There was no funding for the study. The authors have full access to the data and have final responsibility.

References

1. Guzik TJ, Mohiddin SA, Dimarco A, et al. COVID-19 and the cardiovascular system: implications for risk assessment, diagnosis, and treatment options. *Cardiovasc Res*. Aug 1 2020;116(10):1666-1687. doi:10.1093/cvr/cvaa106
2. Wiersinga WJ, Rhodes A, Cheng AC, Peacock SJ, Prescott HC. Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019 (COVID-19): A Review. *JAMA*. Aug 25 2020;324(8):782-793. doi:10.1001/jama.2020.12839
3. Nalbandian A, Sehgal K, Gupta A, et al. Post-acute COVID-19 syndrome. *Nat Med*. Apr 2021;27(4):601-615. doi:10.1038/s41591-021-01283-z
4. Azkur AK, Akdis M, Azkur D, et al. Immune response to SARS-CoV-2 and mechanisms of immunopathological changes in COVID-19. *Allergy*. Jul 2020;75(7):1564-1581. doi:10.1111/all.14364
5. Wang J, Jiang M, Chen X, Montaner LJ. Cytokine storm and leukocyte changes in mild versus severe SARS-CoV-2 infection: Review of 3939 COVID-19 patients in China and emerging pathogenesis and therapy concepts. *Journal of leukocyte biology*. 2020;108(1):17-41. doi:10.1002/JLB.3COVR0520-272R
6. Guo T, Fan Y, Chen M, et al. Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol*. Jul 1 2020;5(7):811-818. doi:10.1001/jamacardio.2020.1017
7. Hendren Nicholas S, Drazner Mark H, Biykem B, Cooper Leslie T. Description and Proposed Management of the Acute COVID-19 Cardiovascular Syndrome. *Circulation*. 2020;141(23):1903-1914. doi:10.1161/CIRCULATIONAHA.120.047349; 27 10.1161/CIRCULATIONAHA.120.047349
8. Yang X, Yu Y, Xu J, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. May 2020;8(5):475-481. doi:10.1016/S2213-2600(20)30079-5
9. Mengozzi A, Georgiopoulos G, Falcone M, et al. The relationship between cardiac injury, inflammation and coagulation in predicting COVID-19 outcome. *Scientific Reports*. 2021/03/22 2021;11(1):6515. doi:10.1038/s41598-021-85646-z
10. Giustino G, Pinney SP, Lala A, et al. Coronavirus and Cardiovascular Disease, Myocardial Injury, and Arrhythmia: JACC Focus Seminar. *Journal of the American College of Cardiology*. 2020;76(17):2011-2023. doi:10.1016/j.jacc.2020.08.059
11. Shi S, Qin M, Shen B, et al. Association of Cardiac Injury With Mortality in Hospitalized Patients With COVID-19 in Wuhan, China. *JAMA Cardiology*. 2020;5(7):802-810. doi:10.1001/jamacardio.2020.0950
12. Bettencourt P, Azevedo A, Pimenta J, Friões F, Ferreira S, Ferreira A. N-Terminal Pro-Brain Natriuretic Peptide Predicts Outcome After Hospital Discharge in Heart Failure Patients. *Circulation*. 2004;110(15):2168-2174. doi:10.1161/01.CIR.0000144310.04433.BE
13. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA*. 2020;323(11):1061-1069. doi:10.1001/jama.2020.1585

14. Argenziano MG, Bruce SL, Slater CL, et al. Characterization and clinical course of 1000 Patients with COVID-19 in New York: retrospective case series. *medRxiv*. Apr 22 2020;doi:10.1101/2020.04.20.20072116
15. Zeng JH, Wu WB, Qu JX, et al. Cardiac manifestations of COVID-19 in Shenzhen, China. *Infection*. Dec 2020;48(6):861-870. doi:10.1007/s15010-020-01473-w
16. Dweck MR, Bularga A, Hahn RT, et al. Global evaluation of echocardiography in patients with COVID-19. *Eur Heart J Cardiovasc Imaging*. Sep 1 2020;21(9):949-958. doi:10.1093/ehjci/jeaa178
17. Jain SS, Liu Q, Raikhelkar J, et al. Indications for and Findings on Transthoracic Echocardiography in COVID-19. *J Am Soc Echocardiogr*. Oct 2020;33(10):1278-1284. doi:10.1016/j.echo.2020.06.009
18. Kang Y, Chen T, Mui D, et al. Cardiovascular manifestations and treatment considerations in COVID-19. *Heart*. Aug 2020;106(15):1132-1141. doi:10.1136/heartjnl-2020-317056
19. Lippi G, Sanchis-Gomar F, Favaloro EJ, Lavie CJ, Henry BM. Coronavirus Disease 2019-Associated Coagulopathy. *Mayo Clin Proc*. Jan 2021;96(1):203-217. doi:10.1016/j.mayocp.2020.10.031
20. Gao L, Jiang D, Wen X-S, et al. Prognostic value of NT-proBNP in patients with severe COVID-19. *Respiratory research*. 2020;21(1):83. doi:10.1186/s12931-020-01352-w
21. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. Feb 15 2020;395(10223):507-513. doi:10.1016/S0140-6736(20)30211-7
22. Clerkin KJ, Fried JA, Raikhelkar J, et al. COVID-19 and Cardiovascular Disease. *Circulation*. May 19 2020;141(20):1648-1655. doi:10.1161/CIRCULATIONAHA.120.046941
23. D F. The cardiovascular impact of COVID-19. *Diagnostic and Interventional Cardiology (DAIC)*. 2020;
24. Covidence systematic review software. Veritas Health Innovation. Melbourne, Australia.
25. National Health Commission of the People's Republic of China. Chinese management guideline for COVID-19 (version 7.0) (2020).
26. World Health O. *Clinical management of severe acute respiratory infection when novel coronavirus (2019-nCoV) infection is suspected: interim guidance, 28 January 2020*. 2020. 2020. <https://apps.who.int/iris/handle/10665/330893>
27. Thygesen K, Alpert JS, Jaffe AS, et al. Fourth universal definition of myocardial infarction (2018). *European Heart Journal*. 2018;40(3):237-269. doi:10.1093/eurheartj/ehy462
28. Wells G SB, O'Connell D, Peterson J, Welch V, Losos M, Tugwell P. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp
29. McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Research Synthesis Methods*. 2020/04/26 2020;n/a(n/a)doi:10.1002/jrsm.1411
30. *Stata: Release 16. Statistical Software*. 2021.
31. Chang BH, Hoaglin DC. Meta-Analysis of Odds Ratios: Current Good Practices. *Med Care*. Apr 2017;55(4):328-335. doi:10.1097/MLR.0000000000000696

32. Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med*. Jun 15 2002;21(11):1539-58. doi:10.1002/sim.1186
33. Lin L, Chu H. Quantifying publication bias in meta-analysis. *Biometrics*. 2018;74(3):785-794. doi:10.1111/biom.12817
34. Nyaga VN, Arbyn M, Aerts M. Metaprop: a Stata command to perform meta-analysis of binomial data. *Arch Public Health*. 2014;72(1):39. doi:10.1186/2049-3258-72-39
35. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *Int J Surg*. Apr 2021;88:105906. doi:10.1016/j.ijsu.2021.105906
36. *PRISMA2020: R package and ShinyApp for producing PRISMA 2020 compliant flow diagrams (Version 0.0.1)*. Zenodo; 2020. <http://doi.org/10.5281/zenodo.4287835>
37. Chung MK, Zidar DA, Bristow MR, et al. COVID-19 and Cardiovascular Disease: From Bench to Bedside. *Circ Res*. Apr 16 2021;128(8):1214-1236. doi:10.1161/CIRCRESAHA.121.317997
38. Khan S, Rasool ST, Ahmed SI. Role of Cardiac Biomarkers in COVID-19: What Recent Investigations Tell Us? *Curr Probl Cardiol*. 2021:100842-100842. doi:10.1016/j.cpcardiol.2021.100842
39. Lippi G, Lavie CJ, Sanchis-Gomar F. Cardiac troponin I in patients with coronavirus disease 2019 (COVID-19): Evidence from a meta-analysis. *Progress in cardiovascular diseases*. 2020;63(3):390-391. doi:10.1016/j.pcad.2020.03.001
40. Ogungbe OC-M, Y. . Cardiac Biomarkers and COVID-19: What Nurses Should Know About Cardiac and Inflammatory Biomarkers and COVID-19. PCNA (Preventive Cardiovascular Nurses Association); 2021. <https://pcna.net/biomarkers-and-covid-19/>
41. Liu PP, Blet A, Smyth D, Li H. The Science Underlying COVID-19: Implications for the Cardiovascular System. (1524-4539 (Electronic))
42. Lindner D, Fitzek A, Brauningner H, et al. Association of Cardiac Infection With SARS-CoV-2 in Confirmed COVID-19 Autopsy Cases. *JAMA Cardiol*. Nov 1 2020;5(11):1281-1285. doi:10.1001/jamacardio.2020.3551
43. Gheblawi M, Wang K, Viveiros A, et al. Response by Gheblawi et al to Letter Regarding Article, "Angiotensin-Converting Enzyme 2: SARS-CoV-2 Receptor and Regulator of the Renin-Angiotensin System: Celebrating the 20th Anniversary of the Discovery of ACE2". *Circ Res*. Jul 3 2020;127(2):e46-e47. doi:10.1161/CIRCRESAHA.120.317332
44. Scott AJ, O'Dea KP, O'Callaghan D, et al. Reactive oxygen species and p38 mitogen-activated protein kinase mediate tumor necrosis factor α -converting enzyme (TACE/ADAM-17) activation in primary human monocytes. *J Biol Chem*. 2011;286(41):35466-35476. doi:10.1074/jbc.M111.277434
45. Mogensen TH. Pathogen recognition and inflammatory signaling in innate immune defenses. (1098-6618 (Electronic))
46. Vallance P, Collier J Fau - Bhagat K, Bhagat K. Infection, inflammation, and infarction: does acute endothelial dysfunction provide a link? (0140-6736 (Print))
47. Jin Y, Ji W, Yang H, Chen S, Zhang W, Duan G. Endothelial activation and dysfunction in COVID-19: from basic mechanisms to potential therapeutic approaches. *Signal Transduction and Targeted Therapy*. 2020/12/24 2020;5(1):293. doi:10.1038/s41392-020-00454-7

48. Acanfora D, Acanfora C, Ciccone MM, et al. The Cross-Talk between Thrombosis and Inflammatory Storm in Acute and Long-COVID-19: Therapeutic Targets and Clinical Cases. *Viruses*. 2021;13(10):1904. doi:10.3390/v13101904
49. Puntmann VO, Carerj ML, Wieters I, et al. Outcomes of Cardiovascular Magnetic Resonance Imaging in Patients Recently Recovered From Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol*. Nov 1 2020;5(11):1265-1273. doi:10.1001/jamacardio.2020.3557
50. Datta SD, Talwar A, Lee JT. A Proposed Framework and Timeline of the Spectrum of Disease Due to SARS-CoV-2 Infection: Illness Beyond Acute Infection and Public Health Implications. *JAMA*. 2020;324(22):2251-2252. doi:10.1001/jama.2020.22717
51. Abdeladim S, Oualim S, Elouarradi A, et al. Analysis of cardiac injury biomarkers in COVID-19 patients. *Archives of Clinical Infectious Diseases*. 2020;15(4):1-5. doi:10.5812/archcid.105515
52. Cao JT, Zheng Y, Luo Z, et al. Myocardial injury and COVID-19: Serum hs-cTnI level in risk stratification and the prediction of 30-day fatality in COVID-19 patients with no prior cardiovascular disease. *Theranostics*. 2020;10(21):9663-9673. doi:10.7150/thno.47980
53. Cao L, Zhang S, Luo X, et al. Myocardium injury biomarkers predict prognosis of critically ill coronavirus disease 2019 (Covid-19) patients. *Ann Palliat Med*. 2020;9(6):4156-4165. doi:10.21037/apm-20-2112
54. Bardají A, Carrasquer A, Sánchez-Giménez R, et al. Prognostic implications of myocardial injury in patients with and without COVID-19 infection treated in a university hospital. *Rev Esp Cardiol (Engl Ed)*. 2020;doi:10.1016/j.rec.2020.08.027
55. Deng P, Ke Z, Ying B, Qiao B, Yuan L. The diagnostic and prognostic role of myocardial injury biomarkers in hospitalized patients with COVID-19. *Clin Chim Acta*. 2020;510:186-190. doi:10.1016/j.cca.2020.07.018
56. Deng Q, Hu B, Zhang Y, et al. Suspected myocardial injury in patients with COVID-19: Evidence from front-line clinical observation in Wuhan, China. *Int J Cardiol*. 2020;311:116-121. doi:10.1016/j.ijcard.2020.03.087
57. Chen LQ, Burdowski J, Marfatia R, et al. Reduced cardiac function is associated with cardiac injury and mortality risk in hospitalized COVID-19 Patients. *Clin Cardiol*. 2020;43(12):1547-1554. doi:10.1002/clc.23479
58. Chen Q, Xu L, Dai Y, et al. Cardiovascular manifestations in severe and critical patients with COVID-19. *Clin Cardiol*. 2020;43(7):796-802. doi:10.1002/clc.23384
59. Doyen D, Dupland P, Morand L, et al. Characteristics of cardiac injury in critically ill patients with COVID-19. *Chest*. 2020;doi:10.1016/j.chest.2020.10.056
60. Fan H, Zhang L, Huang B, et al. Cardiac injuries in patients with coronavirus disease 2019: Not to be ignored. *Int J Infect Dis*. 2020;96:294-297. doi:10.1016/j.ijid.2020.05.024
61. Fan Q, Zhu H, Zhao J, et al. Risk factors for myocardial injury in patients with coronavirus disease 2019 in China. *ESC Heart Fail*. 2020;doi:10.1002/ehf2.13022
62. Ferrante G, Fazzari F, Cozzi O, et al. Risk factors for myocardial injury and death in patients with COVID-19: insights from a cohort study with chest computed tomography. *Cardiovasc Res*. 2020;116(14):2239-2246. doi:10.1093/cvr/cvaa193
63. Gao L, Jiang D, Wen X-s, et al. Prognostic value of NT-proBNP in patients with severe COVID-19. *Respiratory Research*. 2020;21(1):1-7. doi:10.1186/s12931-020-01352-w

64. Ghio S, Baldi E, Vicentini A, et al. Cardiac involvement at presentation in patients hospitalized with COVID-19 and their outcome in a tertiary referral hospital in Northern Italy. *Internal and emergency medicine*. 2020;15(8):1457-1465. doi:10.1007/s11739-020-02493-y
65. Giustino G, Croft LB, Stefanini GG, et al. Characterization of Myocardial Injury in Patients With COVID-19. *Journal of the American College of Cardiology (JACC)*. 2020;76(18):2043-2055. doi:10.1016/j.jacc.2020.08.069
66. Guo H, Shen Y, Wu N, Sun X. Myocardial injury in severe and critical coronavirus disease 2019 patients. *J Card Surg*. 2020;doi:10.1111/jocs.15164
67. Guo T, Fan Y, Chen M, et al. Cardiovascular Implications of Fatal Outcomes of Patients with Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol*. 2020;5(7):811-818. doi:10.1001/jamacardio.2020.101731950516; Hui, D.S., Azhar, E.I., Madani, T.A., The continuing 2019-nCoV epidemic threat of novel coronaviruses to global health: The latest 2019 novel coronavirus outbreak in Wuhan, China (2020) *Int J Infect Dis*, 91, pp. 264-2
68. Lala A, Johnson KW, Januzzi JL, et al. Prevalence and Impact of Myocardial Injury in Patients Hospitalized With COVID-19 Infection. *Journal of the American College of Cardiology (JACC)*. 2020;76(5):533-546. doi:10.1016/j.jacc.2020.06.007
69. Han H, Xie L, Liu R, et al. Analysis of heart injury laboratory parameters in 273 COVID-19 patients in one hospital in Wuhan, China. *J Med Virol*. 2020;92(7):819-823. doi:10.1002/jmv.25809
70. He X, Wang L, Wang H, et al. Factors associated with acute cardiac injury and their effects on mortality in patients with COVID-19. *Sci Rep*. 2020;10(1)doi:10.1038/s41598-020-77172-1
71. Heberto AB, Carlos PCJ, Antonio CRJ, et al. Implications of myocardial injury in Mexican hospitalized patients with coronavirus disease 2019 (COVID-19). *Int J Cardiol Heart Vasc*. 2020;30:100638. doi:10.1016/j.ijcha.2020.100638
72. Jin L, Tang W, Song L, et al. Acute cardiac injury in adult hospitalized COVID-19 patients in Zhuhai, China. *Cardiovasc Diagn Theer*. 2020;10(5):1303-1312. doi:10.21037/cdt-20-607
73. Almeida Junior GLG, Braga F, Jorge JK, et al. Prognostic Value of Troponin-T and B-Type Natriuretic Peptide in Patients Hospitalized for COVID-19. *Arq Bras Cardiol*. 2020;115(4):660-666. doi:10.36660/abc.20200385
74. Maeda T, Obata R, Rizk D, Kuno T. Cardiac Injury and Outcomes of Patients with COVID-19 in New York City. *Heart Lung Circul*. 2020;doi:10.1016/j.hlc.2020.10.025
75. Majure DT, Gruberg L, Saba SG, Kvasnovsky C, Hirsch JS, Jauhar R. Usefulness of Elevated Troponin to Predict Death in Patients With COVID-19 and Myocardial Injury. *American Journal of Cardiology*. 2020;doi:10.1016/j.amjcard.2020.09.060
76. Kim IC, Song JE, Lee HJ, et al. The Implication of Cardiac Injury Score on In-hospital Mortality of Coronavirus Disease 2019. *J Korean Med Sci*. 2020;35(39):1/10-10/10. doi:10.3346/jkms.2020.35.e349
77. Li C, Jiang J, Wang F, et al. Longitudinal correlation of biomarkers of cardiac injury, inflammation, and coagulation to outcome in hospitalized COVID-19 patients. *J Mol Cell Cardiol*. 2020;147:74-87. doi:10.1016/j.yjmcc.2020.08.008
78. Li J, Zhang Y, Wang F, et al. Cardiac damage in patients with the severe type of coronavirus disease 2019 (COVID-19). *BMC Cardiovascular Disorders*. 2020;20(1):N.PAG-N.PAG. doi:10.1186/s12872-020-01758-w

79. Lombardi CM, Carubelli V, Iorio A, et al. Association of Troponin Levels with Mortality in Italian Patients Hospitalized with Coronavirus Disease 2019: Results of a Multicenter Study. *JAMA Cardiol.* 2020;5(11):1274-1280. doi:10.1001/jamacardio.2020.353831389986; Shi, S., Qin, M., Shen, B., Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China (2020) *Jama Cardiol.*,
<http://jamanetwork.com/article.aspx?doi=10.1001/jamacardio.2020.09>
80. Metkus TS, Sokoll LJ, Barth AS, et al. Myocardial Injury in Severe COVID-19 Compared to Non-COVID Acute Respiratory Distress Syndrome. *Circulation.* 2020;doi:10.1161/CIRCULATIONAHA.120.050543
81. Mu S, Wei W, Jin C, et al. Risk factors for COVID-19 patients with cardiac injury: pulmonary ventilation dysfunction and oxygen inhalation insufficiency are not the direct causes. *Aging.* 2020;12doi:10.18632/aging.104148
82. Karbalai Saleh S, Oraii A, Soleimani A, et al. The association between cardiac injury and outcomes in hospitalized patients with COVID-19. *Intern Emerg Med.* 2020;15(8):1415-1424. doi:10.1007/s11739-020-02466-1
83. Salvatici M, Barbieri B, Cioffi SMG, et al. Association between cardiac troponin I and mortality in patients with COVID-19. *Biomarkers.* 7. doi:10.1080/1354750x.2020.1831609
84. Schiavone M, Gasperetti A, Mancone M, et al. Redefining the Prognostic Value of High-Sensitivity Troponin in COVID-19 Patients: The Importance of Concomitant Coronary Artery Disease. *J Clin Med.* 2020;9(10):13. doi:10.3390/jcm9103263
85. Qian H, Gao P, Tian R, et al. Myocardial Injury on Admission as a Risk in Critically Ill COVID-19 Patients: A Retrospective in-ICU Study. *J Cardiothorac Vasc Anesth.* 2020;doi:10.1053/j.jvca.2020.10.019
86. Qin J-J, Cheng X, Zhou F, et al. Redefining Cardiac Biomarkers in Predicting Mortality of Inpatients With COVID-19. *Hypertension (0194911X).* 2020;76(4):1104-1112. doi:10.1161/HYPERTENSIONAHA.120.15528
87. Raad M, Dabbagh M, Gorgis S, et al. Cardiac Injury Patterns and Inpatient Outcomes Among Patients Admitted With COVID-19. *Am J Cardiol.* 2020;133:154-161. doi:10.1016/j.amjcard.2020.07.040
88. van den Heuvel FMA, Vos JL, Koop Y, et al. Cardiac function in relation to myocardial injury in hospitalised patients with COVID-19. *Neth Heart J.* 2020;28(7-8):410-417. doi:10.1007/s12471-020-01458-2
89. Wang Y, Zheng Y, Tong Q, et al. Cardiac Injury and Clinical Course of Patients With Coronavirus Disease 2019. *Front Cardiovasc Med.* 2020;7:147. doi:10.3389/fcvm.2020.00147
90. Wei JF, Huang FY, Xiong TY, et al. Acute myocardial injury is common in patients with COVID-19 and impairs their prognosis. *Heart.* 2020;106(15):1154-1159. doi:10.1136/heartjnl-2020-317007
91. Shi S, Qin M, Cai Y, et al. Characteristics and clinical significance of myocardial injury in patients with severe coronavirus disease 2019. *Eur Heart J.* 2020;41(22):2070-2079. doi:10.1093/eurheartj/ehaa408
92. Shi S, Qin M, Shen B, et al. Association of Cardiac Injury with Mortality in Hospitalized Patients with COVID-19 in Wuhan, China. *JAMA Cardiol.* 2020;5(7):802-810. doi:10.1001/jamacardio.2020.095031986264; Wang, D., Hu, B., Hu, C., Clinical characteristics of

- 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China (2020) *Jama*, , <http://jamanetwork.com/article.aspx?doi=10.1001/jama.2020.1>
93. Yan X, Wang S, Ma P, et al. Cardiac injury is associated with inflammation in geriatric COVID-19 patients. *J Clin Lab Anal*. 2020;doi:10.1002/jcla.23654
 94. Yang C, Liu F, Liu W, et al. Myocardial injury and risk factors for mortality in patients with COVID-19 pneumonia. *Int J Cardiol*. 2020;doi:10.1016/j.ijcard.2020.09.048
 95. Yang J, Liao X, Yin W, et al. Elevated cardiac biomarkers may be effective prognostic predictors for patients with COVID-19: A multicenter, observational study. *American Journal of Emergency Medicine*. 2021;39:34-41. doi:10.1016/j.ajem.2020.10.013
 96. Zaninotto M, Mion MM, Padoan A, Babuin L, Plebani M. Cardiac troponin I in SARS-CoV-2-patients: The additional prognostic value of serial monitoring. *Clin Chim Acta*. 2020;511:75-80. doi:10.1016/j.cca.2020.09.036
 97. Fan ZX, Yang J, Zhang J, et al. Analysis of influencing factors related to elevated serum troponin I level for COVID-19 patients in Yichang, China. *Cardiovasc Diagn Ther*. 2020;10(4):678-686. doi:10.21037/cdt-20-510
 98. Zhou W, Song L, Wang X, et al. Cardiac injury prediction and lymphocyte immunity and inflammation analysis in hospitalized patients with coronavirus disease 2019 (COVID-19). *Int J Cardiol*. 2020;doi:10.1016/j.ijcard.2020.10.049
 99. Barman HA, Atici A, Sahin I, et al. Prognostic significance of cardiac injury in COVID-19 patients with and without coronary artery disease. *Coron Artery Dis*. Aug 1 2021;32(5):359-366. doi:10.1097/mca.0000000000000914
 100. Chen H, Li X, Marmar T, et al. Cardiac Troponin I association with critical illness and death risk in 726 seriously ill COVID-19 patients: A retrospective cohort study. *Int J Med Sci*. 2021;18(6):1474-1483. doi:10.7150/ijms.53641
 101. De Marzo V, Di Biagio A, Della Bona R, et al. Prevalence and prognostic value of cardiac troponin in elderly patients hospitalized for COVID-19. *J Geriatr Cardiol*. 2021;18(5):338-345. doi:10.11909/j.issn.1671-5411.2021.05.004
 102. Demir OM, Ryan M, Cirillo C, et al. Impact and Determinants of High-Sensitivity Cardiac Troponin-T Concentration in Patients With COVID-19 Admitted to Critical Care. *Am J Cardiol*. 2021;147:129-136. doi:10.1016/j.amjcard.2021.01.037
 103. García de Guadiana-Romualdo L, Morell-García D, Morales-Indiano C, et al. Characteristics and laboratory findings on admission to the emergency department among 2873 hospitalized patients with COVID-19: the impact of adjusted laboratory tests in multicenter studies. A multicenter study in Spain (BIOCOVID-Spain study). *Scand J Clin Lab Invest*. 2021;81(3):187-193. doi:10.1080/00365513.2021.1881997
 104. He J, Zhang B, Zhou Q, et al. The prognostic value of myocardial injury in COVID-19 patients and associated characteristics. *Immun Inflamm Dis*. 2021;doi:10.1002/iid3.484
 105. Liaqat A, Ali-Khan RS, Asad M, Rafique Z. Evaluation of myocardial injury patterns and ST changes among critical and non-critical patients with coronavirus-19 disease. *Sci Rep*. 2021;11(1):4828. doi:10.1038/s41598-021-84467-4
 106. Lu Y, Huang Z, Wang M, et al. Clinical characteristics and predictors of mortality in young adults with severe COVID-19: a retrospective observational study. *Annals of Clinical Microbiology and Antimicrobials*. 2021;20(1)doi:10.1186/s12941-020-00412-9

107. Maeda T, Obata R, Rizk D, Kuno T. Cardiac Injury and Outcomes of Patients With COVID-19 in New York City. *Heart, Lung & Circulation*. 2021;30(6):848-853. doi:10.1016/j.hlc.2020.10.025
108. Mengozzi A, Georgiopoulos G, Falcone M, et al. The relationship between cardiac injury, inflammation and coagulation in predicting COVID-19 outcome. *Sci Rep*. 2021;11(1):6515. doi:10.1038/s41598-021-85646-z
109. Omar T, Karakayalı M, Perincek G. Assessment of COVID-19 deaths from cardiological perspective. *Acta Cardiol*. 2021:1-8. doi:10.1080/00015385.2021.1903704
110. Selçuk M, Keskin M, Çınar T, et al. Prognostic significance of N-Terminal Pro-BNP in patients with COVID-19 pneumonia without previous history of heart failure. *Journal of Cardiovascular and Thoracic Research*. 2021;13(2):141-145. doi:10.34172/jcvtr.2021.26
111. Wang Y, Shu H, Liu H, et al. The peak levels of highly sensitive troponin I predicts in-hospital mortality in COVID-19 patients with cardiac injury: a retrospective study. *Eur Heart J Acute Cardiovasc Care*. 2021;10(1):6-15. doi:10.1093/ehjacc/zuaa019
112. Zhu F, Li W, Lin Q, Xu M, Du J, Li H. Myoglobin and troponin as prognostic factors in patients with COVID-19 pneumonia. *Med Clin (Barc)*. Feb 27 2021;doi:10.1016/j.medcli.2021.01.013




Figure 1. PRISMA flowchart showing study search and selection

Table 1. Characteristics of studies examining cardiac biomarkers and COVID-19 outcomes, N=62

| First Author, Year | Country | Sample size | Sex | | Study Design | Mean Age, (±SD); Median Age (IQR) | Population | Main Cardiovascular Outcome/biomarker assessed | Primary Outcome assessed |
|------------------------------|--------------|-------------|---------------|-------------|---------------|-----------------------------------|--|---|---|
| | | | Female, n (%) | Male, n (%) | | | | | |
| Abdeladim 2020 ⁵¹ | Morocco | 73 | 45 (61.6) | 28 (38.4) | Retrospective | 54.6 (±17.6) | Hospitalized COVID-19 patients with Mild, Severe and Critical cases | Myocardial Injury (cTnI) | Death, severe COVID-19 |
| Cao I 2020 ⁵² | China | 244 | 111 (45.5) | - | Retrospective | 62.6 (±13.4) | Hospitalized COVID-19 patients with no preexisting cardiovascular disease or renal dysfunction | Myocardial Injury (hs-cTnI) | Death (Non-survivors) |
| Cao II 2020 ⁵³ | China | 100 | 55 (55) | 45 (45) | Retrospective | 71.1 (±14.5) | Critically ill COVID-19 patients on ICU admission | Myocardial Injury (MYO, CK-MB and hs-cTnI) | Death |
| Bardaji 2020 ⁵⁴ | Spain | 433 | 174 (40.3) | 259 (59.7) | Retrospective | 67.5 (52.5-77.5) | Consecutive hospitalized patients treated for suspected SARS-CoV-2 infection | Myocardial Injury (cTnI) | 30-day mortality |
| Deng I 2020 ⁵⁵ | China | 264 | - | 130 (49.2) | Retrospective | 64.5 (53.3-74.0) | Inpatients who had a COVID-19 diagnosis. | Cardiac biomarkers (cTnI-ultra, CKMB, MYO, NT-proBNP) | In-hospital death |
| Deng II 2020 ⁵⁶ | China | 112 | 55 (49.1) | 57 (50.9) | Retrospective | 65 (49-70.8) | Diagnosed hospitalized COVID-19 Patients | Myocardial injury, myocarditis ^a and cardiac dysfunction | Composite endpoint of ICU admission, MV, ECMO, or death |
| Chen I 2020 ⁵⁷ | US | 143 | - | 89 (62) | Retrospective | 67 (±16) | Laboratory confirmed hospitalized COVID-19 patients | Cardiac function (cTnI & Echocardiograms) | In-hospital death |
| Chen II 2020 ⁵⁸ | China | 54 | 18 (33.3) | 36 (66.7) | Retrospective | 57.6 (44.9-70.3) | Hospitalized COVID-19 patients | Myocardial Injury (TnI) | Severe COVID-19 (ARDS, Pneumonia) |
| Doyen 2020 ⁵⁹ | France | 43 | 7 (16) | 36 (84) | Prospective | 60 (±13) | COVID-19 patients on ICU admission | Cardiac functioning (ECG, hs-TnI, TTE) | ICU discharge or death |
| Fan I 2020 ⁶⁰ | China | 73 | 24 (32.9) | 49 (67.1) | Retrospective | 58.4 (±14.3) | ICU admitted COVID-19 patients | Myocardial Injury (cTnI) | Death |
| Fan II 2020 ⁶¹ | China | 353 | - | 186 (52.7) | Retrospective | 60.5 (±10.1) | Hospitalized patients with confirmed COVID-19 diagnosis, | Myocardial Injury (cTnI) | Severity of myocardial injury |
| Ferrante 2020 ⁶² | Italy | 332 | 95 (28.6) | (71.4) | Retrospective | 66.9 (55.4-75.5) | Patients on ICU admission with COVID-19 | Myocardial Injury (hs-cTnI) | Death, duration of Hospital and admission into ICU |
| Gao 2020 ⁶³ | China | 54 | 30 (55.6) | 24 (44.4) | Retrospective | 60.4 (±16.1) | Patients with severe COVID-19 | In-hospital death | Death |
| Ghio 2020 ⁶⁴ | Italy | 340 | 107 (31.4) | 233 (68.6) | Retrospective | 69.8 (58.6-78.3) | Adult patients admitted with the suspected diagnosis of COVID-19 | Myocardial injury (hs-TnI) | Death and/or ICU admission, MV |
| Giustino 2020 ⁶⁵ | US and Italy | 305 | 100 (32.8) | 205 (67.2) | Retrospective | 63 (53-73) | Hospitalized patients with COVID-19 | Myocardial Injury (hs-TnI), TTE abnormalities | Death |
| Guo I 2020 ⁶⁶ | China | 74 | 25 (33.8) | 49 (66.2) | Retrospective | 67.2 (±14.6) | Patients on ICU admission with severe or critical COVID-19 | Myocardial Injury (cTnI, NT-proBNP, CK-MB, MYO) | Death, MV time |

| | | | | | | | | | |
|-----------------------------------|--------|--------|--------------|--------------|---------------|-------------------|--|---|--|
| Guo II 2020 ⁶⁷ | China | 187 | - | 91(48.7) | Retrospective | 58.5 (±14.7) | Hospitalized patients with confirmed COVID-19 | Myocardial Injury (TnT) | Death |
| Lala 2020 ⁶⁸ | US | 2736 | 1,106 (40.4) | 1630 (59.6) | Retrospective | 66.4 | Patients admitted with a laboratory-confirmed SARS-CoV-2 infection | Myocardial Injury (TnI) | Death, intubation, or hospital discharge |
| Han 2020 ⁶⁹ | China | 273 | 176 (64.5) | 97 (35.53) | Retrospective | 58.9 (±10.8) | Hospitalized patients diagnosed with SARS-CoV-2 infection | Cardiac biomarkers (ultra-TnI, NT-proBNP, CK-MB, MYO) | COVID-19 severity |
| He I 2020 ⁷⁰ | China | 1031 | 493 (47.8) | 538 (52.2) | Retrospective | 63 (52-70) | Inpatients with laboratory-confirmed COVID-19 | Acute cardiac Injury (hs-TnI) | Severe COVID-19 (ARDS, Pneumonia) |
| Heberto 2020 ⁷¹ | Mexico | 254 | 87 (34.3) | 167 (65.7) | Prospective | 53.8 (±12.7) | Hospitalized patients, with and without myocardial injury, and a confirmed test result of COVID-19 | Cardiac markers (hs-TnI, NT-proBNP) | COVID-19 complications: MV, Death |
| Jin 2020 ⁷² | China | 93 | 52 (55.9) | 41 (44.1) | Retrospective | 48.0 (35.5-62.5) | COVID-19 patients on hospital admission | Myocardial Injury (hs-TnI, CK-MB, CMR) | - |
| Almeida Junior 2020 ⁷³ | Brazil | 183 | 63 (34.4) | 120 (65.59) | Retrospective | 66.8 (±17) | Patients admitted for COVID-19 | Myocardial Injury (TnT-ultra, BNP) | ICU admission, MV, all-cause death |
| Maeda I 2020 ⁷⁴ | US | 224 | - | 125 (55.8) | Retrospective | 64.0 (±16.6) | Patients admitted with COVID-19 | Myocardial Injury (cTnI), ECG abnormality | In-hospital death |
| Majure 2020 ⁷⁵ | US | 11,159 | 4464 (40) | 6695 (60) | Retrospective | 66 (56-77) | Hospitalized patients diagnosed with COVID 19 | Myocardial Injury (cTnI, TnT), ECG abnormality | In-hospital death |
| Kim 2020 ⁷⁶ | Korea | 38 | 16 (40.9) | 22 (59.1) | Retrospective | 69.6 (±14.9) | In-patients with confirmed COVID-19 | Myocardial Injury (cTnI, CK-MB) | - |
| Li I 2020 ⁷⁷ | China | 2,068 | - | 1,005 (48.6) | Retrospective | 63 (51-70) | Critical and non-critical hospitalized COVID-19 patients | Myocardial Injury (hs-cTnI) | Death |
| Li II 2020 ⁷⁸ | China | 100 | 56 (56) | 44 (44) | Retrospective | 62.0 (51.0-70.8) | Patients admitted with COVID-19 | Myocardial Injury (hs-cTnI, CK-MB, MYO) | Discharge, death |
| Lombardi 2020 ⁷⁹ | Italy | 614 | 179 (29.2) | 435 (70.8) | Retrospective | 67 (±13) | Hospitalized patients with laboratory-confirmed SARS-CoV-2 infection | Myocardial Injury (hs-TnI, TnT) | Discharge, death |
| Metkus 2020 ⁸⁰ | US | 243 | 95 (39.1) | 148 (60.9) | Retrospective | 62.8 (±14.9) | Intubated confirmed COVID-19 patients who underwent troponin assessment 24hrs post intubation | Myocardial Injury (hs-TnI, TnT) | ARDS unrelated to COVID |
| Mu 2020 ⁸¹ | China | 113 | 51 (45.1) | 62 (54.9) | Retrospective | 63.00 (49.5-70.0) | Hospitalized patients with confirmed COVID-19 diagnosis | Myocardial Injury (cTnI) | Severe COVID-19 (ARDS, Pneumonia) |
| Karbalai Saleh 2020 ⁸² | Iran | 386 | 58 (38.9) | - | Retrospective | 59.5 (±15.8) | Hospitalized patients with COVID-19 | Myocardial Injury (hs-TnI), ECG abnormalities | Death, ICU admission |
| Salvatici, 2020 ⁸³ | Italy | 1,055 | 339 (32.1) | 355 (67.9) | Retrospective | 64.4 ±14.2 | Hospitalized patients with confirmed SARS-CoV-2 | Myocardial Injury (cTnI) | Death |
| Schiavone 2020 ⁸⁴ | Italy | 674 | 268 (39.8) | 406 (60.2) | Retrospective | 60.8 (±15.9) | Hospitalized COVID-19 patients, with or without chronic coronary syndromes | Myocardial Injury (cTnI) | Death |
| Qian 2020 ⁸⁵ | China | 77 | 24 (31.2) | 53 (68.8) | Retrospective | 65.5 (±12.2) | Newly admitted ICU patients with COVID-19 | Myocardial Injury (cTnI) | ICU admission, Death |
| Qin 2020 ⁸⁶ | China | 6,033 | 3174 (52.61) | 2859 (47.39) | Retrospective | 57 (45-66) | Hospitalized COVID-19 patients | Myocardial Injury (hs-TnI, CK-MB) | 28-day mortality, ICU admission, MV |
| Raad 2020 ⁸⁷ | US | 1,020 | 511 (50) | 509 (50) | Retrospective | 63 (52-73) | Hospitalized adult patients diagnosed with SARS-CoV-2 | Myocardial Injury (hs-TnI) | Death and LOS |

| | | | | | | | | | |
|---------------------------------------|-------------|------|------------|-------------|---------------|-------------------------|--|--|--------------------------|
| van den Heuvel 2020 ⁸⁸ | Netherlands | 51 | 10 (20) | 41 (80) | Prospective | 63 (51-68) | Hospitalized patients with COVID-19 | Myocardial Injury (hs-TnI), ventricular dysfunction (TTE) | ICU admission |
| Wang I 2020 ⁸⁹ | China | 222 | 109 (49.1) | 113 (50.9) | Retrospective | 63.0 (50.0-69.0) | Hospitalized patients with laboratory-confirmed COVID-19 infection | Myocardial Injury (hs-TnI) | Death, recovery |
| Wei 2020 ⁹⁰ | China | 101 | - | 54 (53.5) | Prospective | 49 (34-62) | Hospitalized patients with laboratory confirmed SARS-CoV-2 infection | Myocardial Injury (hs-TnI) | ICU admission, MV, death |
| Shi I 2020 ⁹¹ | China | 671 | 349 (52) | 322 (48) | Retrospective | 63 (50-72) | Hospitalized patients with laboratory-confirmed COVID-19 | Myocardial Injury (cTnI) | Death |
| Shi II 2020 ⁹² | China | 416 | 211 (50.7) | - | Retrospective | 64 (21-95) | Hospitalized patients with laboratory-confirmed COVID-19 | Myocardial Injury (hs-TnI, CK-MB, MYO) | Death |
| Yan 2020 ⁹³ | China | 119 | 66 (55.5) | 52 (44.5) | Retrospective | 69 (66-76) | Hospitalized patients over 60years diagnosed with COVID-19 | Cardiac markers (hs-TnI, CK-MB, MYO) | - |
| Yang I 2021 ⁹⁴ | China | 357 | - | 185 (51.8) | Prospective | 56.0 (43.0-68.0) | Hospitalized patients with COVID-19 | Cardiac markers (hs-TnI, CK-MB, MYO) | Death |
| Yang II 2020 ⁹⁵ | China | 203 | 88(43.3) | 115 (56.7) | Retrospective | 62.0 (49.0-69.0) | Hospitalized patients with COVID-19 | Myocardial Injury (TnI) | Death |
| Zaninotto 2020 ⁹⁶ | Italy | 113 | 33 (29) | 80 (71) | Retrospective | 65 (53-75) | Hospitalized patients with confirmed COVID-19 | Myocardial Injury (hs-TnI) | - |
| Fan III 2020 ⁹⁷ | China | 89 | - | 49 (55.1) | Retrospective | 61.8 (±16.1) | Hospitalized patients diagnosed with COVID-19 | Myocardial Injury (TnI) | - |
| Zhou 2020 ⁹⁸ | China | 68 | 34 (50) | 34 (50) | Retrospective | 67 (30-86) | Hospitalized patients diagnosed with COVID-19 | Myocardial Injury (hs-TnI) | Death |
| Barman 2020 ⁹⁹ | Turkey | 607 | - | 334 (55) | Retrospective | 62.5 (±14.3) | Patients hospitalized for COVID-19 | Myocardial Injury (hs-TnI) | Death |
| Chen III, 2021 ¹⁰⁰ | China | 726 | - | 392 (54.1) | Retrospective | 68 (58-77) [§] | Patients hospitalized for COVID-19 | Myocardial Injury (hs-TnI) | Death, severe COVID-19 |
| De Marzo 2021 ¹⁰¹ | Italy | 343 | - | 119 (34.7) | Retrospective | 75 (68-83) | Patients ≥60years hospitalized with COVID-19 | Myocardial Injury (cTnI) | Death |
| Demir 2021 ¹⁰² | UK | 277 | - | 197 (71) | Retrospective | 55.1 (±13.9) | Hospitalized patients with confirmed COVID-19 | Myocardial Injury (hs-TnI) | Death |
| Guadiana-Romualdo 2021 ¹⁰³ | Spain | 2873 | - | 1699 (59.1) | Retrospective | 66 (54-76) | Hospitalized patients with confirmed COVID-19 | - | All-cause mortality |
| He II 2021 ¹⁰⁴ | China | 304 | - | 160 (52.6) | Retrospective | 65 (54-74) | Hospitalized patients diagnosed with COVID-19 | Myocardial Injury (hs-TnI) | Death |
| Liaqat 2021 ¹⁰⁵ | Pakistan | 201 | 82 (40.8) | 119 (59.2) | Retrospective | 44.6 (±15.2) | Hospitalized patients diagnosed with COVID-19 | Myocardial Injury (TnI, CK-MB), Ventricular dysfunction (TTE), ECG abnormalities | COVID-19 severity, death |
| Lu 2021 ¹⁰⁶ | China | 77 | - | 50 (65) | Retrospective | 59 (54-63) | Hospitalized patients diagnosed with COVID-19 | Myocardial Injury (hs-TnI) | Death, COVID-19 severity |
| Maeda II 2021 ¹⁰⁷ | US | 181 | - | 100 (55.8) | Retrospective | 64.0 (±16.6) | Patients admitted with COVID-19 | Myocardial Injury (hs-TnI), ECG abnormalities | In-hospital death |

| | | | | | | | | | |
|------------------------------|--------|-----|---|------------|---------------|--------------|---|---|--|
| Mengozzi 2021 ¹⁰⁸ | Italy | 266 | - | 180 (67.7) | Prospective | 63 (±15) | Patients hospitalized for COVID-19 | Myocardial Injury (hs-TnT) | ARD, MV, ICU admissionIn-hospital, all-cause death |
| Omar 2021 ¹⁰⁹ | Turkey | 355 | - | 172 (48.5) | Retrospective | 58 (35.5-71) | Patients hospitalized for COVID-19 | Myocardial Injury (hs-TnT), ECG abnormalities | Death |
| Selçuk 2021 ¹¹⁰ | Turkey | 137 | - | 72 (52.5) | Retrospective | 55 (±14) | Patients hospitalized for COVID-19 | Cardiac markers (cTnI, NT-proBNP) | In-hospital death |
| Wang II 2021 ¹¹¹ | China | 242 | - | 151 (62.4) | Retrospective | 68 (61-75) | Patients hospitalized for COVID-19 | Myocardial Injury (cTnI) | ARDS, Death |
| Zhu 2021 ¹¹² | China | 499 | - | 253 (50) | Retrospective | 59 (±15) | Hospitalized severe/critically ill patients with COVID-19 | Myocardial Injury (cTnI, MYO) | Death |

ICU: Intensive Care Unit; MV: Mechanical Ventilation; ECMO: extracorporeal membrane oxygenation; LOS: Length of Stay; hs-cTnI: high-sensitivity cardiac troponin I; TnT; Troponin T; NT-proBNP: N-terminal pro-B-type natriuretic peptide; cTnI-ultra: cardiac troponin I-ultra; CK-MB: creatinine kinase-myocardial band; MYO: Myoglobin; CKMB: Creatinine kinase–myocardial band; ECG: Electrocardiogram; TTE: Transtheoracic Echocardiography; CMR: cardiac magnetic resonance.

^aMyocarditis related abnormalities define as: triple elevation in hypersensitive cardiac Troponin I (over 0.12 ng/mL) plus abnormalities on echocardiography and/or electrocardiogram; [§]Critical patients; [†]With cardiac injury.

Figure 2. Pooled proportion for history of hypertension







Figure 3. Association between biomarkers of myocardial injury, stretch and coagulation with severe COVID-19 and death: meta-analysis results





Notes: Hazard ratios and odds ratios with 95% confidence intervals on a logarithmic scale for individual or pooled study data for pair-wise comparison of odds associated with elevated biomarker levels. Logarithmic OR and HR exponentiated for interpretation, for instance, pooled odds ratio for association between troponin elevation and death/severe disease $e^{(logx)}$: $e^{(0.57[95\%CI:0.35 - 0.78])} = OR: 1.77$ (95%CI: 1.42-2.18).



Highlights

- **In** hospitalized COVID-19 patients, there was a high prevalence of underlying comorbidities.
- Elevated biomarkers of subclinical myocardial injury were significantly associated with severe COVID-19 and death.
- Recovered COVID-19 patients may benefit from minimally invasive assessment for myocardial injury markers.
- There is evidence of persisting symptoms suggestive of complications in patients recovered from COVID-19